

Medication Guide
PLEGRIDY® (PLEGG-rih-dee)
(peginterferon beta-1a) injection for subcutaneous use

Read this Medication Guide before you start using PLEGRIDY, and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is the most important information I should know about PLEGRIDY?

PLEGRIDY can cause serious side effects, including:

- Liver problems or worsening of liver problems, including liver failure and death. Symptoms may include: yellowing of your skin or the white part of your eye, nausea, loss of appetite, tiredness, bleeding more easily than normal, confusion, sleepiness, dark colored urine, and pale stools.
- During your treatment with PLEGRIDY you will need to see your healthcare provider and have regular blood tests to check for these possible side effects.
- Depression or suicidal thoughts. Symptoms may include: new or worsening depression (feeling hopeless or bad about yourself), thoughts of hurting yourself or suicide, irritability (getting upset easily), nervousness, or new or worsening anxiety.

Call your healthcare provider right away if you have any of the symptoms listed above.

What is PLEGRIDY?

- PLEGRIDY is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.
- It is not known if PLEGRIDY is safe and effective in people under 18 or over 65 years of age.

Do not take PLEGRIDY if you:

- are allergic to interferon beta or peginterferon, or any of the other ingredients in PLEGRIDY. See the end of this Medication Guide for a complete list of ingredients in PLEGRIDY.

Before using PLEGRIDY, tell your healthcare provider about all of your medical conditions, including if you:

- are being treated for a mental illness or had treatment in the past for any mental illness, including depression and suicidal behavior.
- have or had liver problems.
- have or had low blood cell counts.
- have or had bleeding problems.
- have or had heart problems.
- have or had seizures (epilepsy).
- have or had thyroid problems.
- have or had any kind of autoimmune disease (where the body's immune system attacks the body's own cells).
- are pregnant or plan to become pregnant. It is not known if PLEGRIDY can harm your unborn baby.
 - If you become pregnant while taking PLEGRIDY, talk to your healthcare provider about enrolling in the PLEGRIDY Pregnancy Registry. You can enroll in this registry by calling 1-866-810-1462 or visiting <https://www.plegridypregnancyregistry.com/>. The purpose of this registry is to collect information about the safety of Plegridy during pregnancy.
- are breastfeeding or plan to breastfeed. PLEGRIDY may pass into your breastmilk. Talk to your healthcare provider about the best way to feed your baby if you use PLEGRIDY.

Tell your healthcare provider about all the medicines you take, including prescription and over the counter medicines, vitamins, and herbal supplements.

How should I use PLEGRIDY?

- See the detailed **Instructions for Use** for instructions on how to prepare and inject your dose of PLEGRIDY.
- Use PLEGRIDY exactly as your healthcare provider tells you. A healthcare provider should show you how to

inject your PLEGRIDY before you use it for the first time.

- Your healthcare provider will tell you how much PLEGRIDY to inject and how often to inject PLEGRIDY. Do not inject more than your healthcare provider tells you to.
- When you use PLEGRIDY for the first time, your healthcare provider may tell you to slowly increase your dose.
- PLEGRIDY is given by injection under the skin (subcutaneous injection) of your stomach (abdomen), back of upper arm, or thigh 1 time every 14 days.
- Change (rotate) the site you choose with each injection to help decrease the chance that you will have an injection site reaction. **Do not** inject into an area of the body where the skin is irritated, reddened, bruised, infected, or scarred in any way.
- After 2 hours check your injection site for redness, swelling or tenderness. If you have a skin reaction and it does not clear up in a few days, contact your healthcare provider.
- Always use a new, PLEGRIDY prefilled pen or new, unopened single dose prefilled syringe for each injection.

What are the possible side effects of PLEGRIDY?

PLEGRIDY may cause serious side effects, including:

- **See “What is the most important information I should know about PLEGRIDY?”**
- **seizures.** Some people have had seizures while taking PLEGRIDY, including people who have never had seizures before.
- **serious allergic reactions.** Serious allergic reactions can happen if you take PLEGRIDY. Symptoms may include: itching, swelling of the face, eyes, lips, tongue, or throat, trouble breathing, feeling faint, anxiousness, skin rash, hives, skin bumps. Get emergency help right away if you have any of these symptoms. Talk to your healthcare provider before taking another dose of PLEGRIDY
- **injection site reactions.** PLEGRIDY may commonly cause redness, pain, or swelling at the place where your injection was given. Call your healthcare provider right away if an injection site becomes swollen and painful or the area looks infected and it does not heal within a few days. You may have a skin infection or an area of severe skin damage (necrosis) requiring treatment by a healthcare provider.
- **heart problems, including congestive heart failure.** Call your healthcare provider right away if you have worsening symptoms of heart failure such as shortness of breath or swelling of your lower legs or feet while using PLEGRIDY.
 - Some people using PLEGRIDY may have other heart problems, including low blood pressure, fast or abnormal heart beat, chest pain, heart attack, or a heart muscle problem (cardiomyopathy).
- **blood problems and changes in your blood tests.** PLEGRIDY can decrease your white blood cells or platelets, which can cause an increased risk of infection, bleeding, or anemia and can cause changes in your liver function tests. Your healthcare provider will do tests to monitor for side effects while you use PLEGRIDY.
- **thrombotic microangiopathy (TMA).** TMA is a condition that involves injury to the smallest blood vessels in your body. TMA can also cause injury to your red blood cells (the cells that carry oxygen to your organs and tissues) and your platelets (cells that help your blood clot) and can sometimes lead to death. Your healthcare provider may tell you to stop taking PLEGRIDY if you develop TMA.
- **autoimmune diseases.** Problems with easy bleeding or bruising (idiopathic thrombocytopenia), thyroid gland problems (hyperthyroidism and hypothyroidism), and autoimmune hepatitis have happened in some people who use interferon beta.

The most common side effects of PLEGRIDY include:

- **flu-like symptoms.** Many people who use PLEGRIDY have flu-like symptoms, especially early in the course of therapy. These symptoms are not really the flu. You cannot pass it on to anyone else.
 - You may be able to manage these flu-like symptoms by taking over-the-counter pain and fever reducers and drinking plenty of water.

Flu like symptoms or other common side effects OF PLEGRIDY may include: headache, muscle and joint aches, fever, chills, or tiredness.

Talk with your healthcare provider about ways to help if you develop flu-like symptoms while taking PLEGRIDY.

These are not all of the possible side effects of PLEGRIDY.
Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General Information about the safe and effective use of PLEGRIDY.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use PLEGRIDY for a condition for which it was not prescribed. Do not give PLEGRIDY to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about PLEGRIDY that is written for health professionals.

What are the ingredients in PLEGRIDY?

Active ingredient: peginterferon beta-1a

Inactive ingredients:

- **Single-dose Prefilled Pen:** arginine hydrochloride, glacial acetic acid, polysorbate 20, sodium acetate trihydrate in sterile water for injection
- **Single-dose Prefilled Syringe:** arginine hydrochloride, glacial acetic acid, polysorbate 20, sodium acetate trihydrate in sterile water for injection

For more information, go to www.plegridy.com or call 1-800-456-2255

Manufactured by: Biogen Inc., Cambridge, MA 02142, U.S. License # 1697

PLEGRIDY is a registered trademark of Biogen. ©2013-2020 Biogen

This Medication Guide has been approved by the U.S. Food and Drug Administration

Issued: 03/2020